

Remarks on the species *Dendrobaena jeanneli* Pop, 1948 and its proposed synonymy with *Octolasion cyaneum* (Savigny, 1826) (Oligochaeta: Lumbricidae)

CS. CSUZDI¹ and V.V. POP²

Abstract. The type specimen of the dubious species *Dendrobaena jeanneli* Pop, 1948 was found in the Zoological Museum, Babes-Bolyai University, Cluj. An investigation of the somewhat softened and fragmented specimen allowed us to recognise its most important taxonomic characters such as the position of the clitellum and tubercles, the setal arrangement, the structure of the calciferous glands and the nephridial bladders. According to these characters this species belongs to the genus *Octolasion* Örley, 1885, and on the position of the clitellum and tubercles it is clearly identical with *O. cyaneum* (Savigny, 1826).

The earthworm species *Dendrobaena jeanneli* was described by Pop (1948) from a cave material collected by René Jeannel in Grotte du Rochefort, France. Since the first description it has not been recollected although this species' name regularly occurs in the earthworm literature. Bouché (1972) mentions this species as valid in his book on the French earthworms, also Mršić (1991) and Blakemore (2005) cite this name among the *Dendrobaena* species. The last checklist in which this species name occurs is by Qiu & Bouché (1998).

During a search for the type species of *Allolobophora dugesi* var. *getica* Pop, 1947 in the Zoological Museum of the Babes-Bolyai University, Cluj we happened to find the type specimen of *Dendrobaena jeanneli* as well. Eventhough the specimen is not well preserved, its condition enabled us to give a short redescription

Dendrobaena jeanneli Pop, 1948

Dendrobaena jeanneli Pop, 1948: 15., Pop 1968: 244., Bouché 1972: 404., Zicsi 1982: 443., Blakemore 2005: 25., Mršić 1991: 550., Qiu & Bouché 1998: 194.

Material examined. *Dendrobaena jeanneli*
Typus. CZMB No. 589 (Grotte de la Rochefort,

Dépt. De la Mayenne, comm. De Thorigné, France. Leg. R. Jeannel, 30. XI. 1912. The specimen is softened and broken in two parts just at the clitellum.

Diagnosis. External – Body length cca 100 mm, diameter 3-4 mm. Colour brown (due to the preservation?), but pigmentation seems to be lacking. Head epilobous, dorsal pores cannot be found. Setae in closer pairs laterally and somewhat wider ventrally; setal arrangement after the clitellum: aa:ab:bc:cd:dd = 4.2:1.8:2:1:11 Clitellum extends on segments 29–34, saddle-shaped. Tubercles as white bands on $\frac{1}{2}29$ – $\frac{1}{2}34$. Male pore on 15 between setae *b*–*c*, great, usually intruding into the neighbouring segments. Nephropores not seen. *Internal* – Septa clearly seen but not tickened. Four pairs of seminal vesicles in 9–12. Spermathecae two pairs in 9/10, 10/11, open in setal lines *cd*. Calciferous glands in 10–14, with well-detached lateral diverticula in 10. Last pair of hearts in 11 and a pair of extraoesophageals in 12. Excretory system holoic. Nephridial bladders ocarina-shaped.

Remarks. Perhaps the darker brown colour of the specimen misled Pop (1948) in identifying its real generic placement. However, due to the shape of

¹Dr. Csaba Csuzdi, Systematic Zoology Research Group of Hungarian Academy of Sciences, and Department of Zoology, Hungarian Natural History Museum, 1088 Budapest, Baross str. 13, Hungary. E-mail: csuzdi01@elte.hu.

²Dr. Victor V. Pop, Institute of Biological Research, 48 Republicii Street. POBox 229, 400015 Cluj-Napoca, Romania, E-mail: victorpop@yahoo.com.

nephridial bladders and the structures of calciferous glands this species could not belong to *Dendrobaena* but to *Octolasion*. On the position of the clitellum and tubercles, this specimen is clearly identical with *O. cyaneum*, therefore *Dendrobaena jeanneli* Pop, 1948 must be regarded as synonym of *Octolasion cyaneum* (Savigny, 1826).

Acknowledgement – This study was partly supported by the Hungarian Scientific Research Grant (No. T42745), and by the Romanian Scientific Grant CEEEx No. 05-D11-82.

REFERENCES

- BLAKEMORE, R. (2005): *An updated list of valid, invalid and synonymous names of Criodriloida and Lumbricoidea (Annelida: Oligochaeta: Criodrilidae, Sparganophilidae, Ailoscolecidae, Hormogastridae, Lumbricidae, Lutodrilidae)*. – In: M.T. ITO and N. KANEKO (eds.): *A Series of Searchable Texts on Earthworm Biodiversity, Ecology and Systematics from Various Regions of the World*. CD edition, Yokohama University, 68 pp.
- BOUCHÉ, M. B. (1972): *Lombriciens de France. Écologie et Systématique*. – Institut National de la Recherche Agronomique, Articles de Zoologie-Écologie animale (Numéro hors-série), 671 pp.
- MRŠIĆ, N. (1991): *Monograph on earthworms (Lumbricidae) of the Balkans I-II*. - Slovenska Akademija Znanosti in Umetnosti, Ljubljana, Opera 31, 757 pp.
- QIU, J-P. & BOUCHÉ, M.B. (2000): Liste classée des taxons valides de Lombriciens (Oligochaeta: Lumbricoidea) après l'étude des trios cinquième d'entre-eux. – *Documents pedozoologiques & integrologiques*, 4: 181-200.
- POP, V. (1948): *Dendrobaena jeanneli*, nouvelle espèce de Lumbricide cavernicole. *Comptes Rendus des Séances du Cercle zoologique de Cluj*, 1947-1848: 15-16.
- POP, V. (1968): Les Lumbricidés cavernicoles de la Collection Biospeologica. – *Archives de Zoologie expérimentale et générale*, 109(2): 229-256.
- ZICSI, A. (1982): Verzeichnis der bis 1971 beschriebenen und revidierten Taxa der Familie Lumbricidae (Oligochaeta). *Acta zoologica hungarica*, 28: 421-454.

INDEX

ANDRÁSSY, I.: Contributions to the genus <i>Opisthodorylaimus</i> Ahmad & Jairajpuri, 1982 (Nematoda: Dorylaimida), with descriptions of two new species.....	3
BLAKEMORE R. J., CSUZDI, CS., ITO, M. T., KANEKO, N. M., PAOLETTI, G., SPIRIDONOV, S. E. UCHIDA, T. & VAN PRAAGH, B. D.: <i>Megascolex (Promegascolex) mekongianus</i> Cognetti, 1922 - its extent, ecology and allocation to <i>Amyntas</i> (Clitellata/Oligochaeta: Megascolecidae).....	19
BOROS, G.: The enchytraeid fauna (Annelida: Oligochaeta) of the Sas-hegy Nature Conservation Area, Hungary	31
KONTSCHÁN, J.: New records for the Uropodina fauna of Bulgaria with descriptions of two new species (Acari: Uropodidae)	37
KONTSCHÁN, J.: Trachyuropodid mites of the Carpathian Basin (Acari Uropodina: Trachyuropodidae).....	43
MAHUNKA, S.: Oribatids from the Carpathian Basin with zoogeographical and taxonomical notes (Acari: Oribatida), II	57
MAHUNKA, S. & MAHUNKA-PAPP, L.: Contribution to the knowledge of the Hungarian Oribatida fauna (Acari) II	69
NÉDLI, J., FORRÓ, L., KORPONAI, J. & G-TÓTH, L.: <i>Daphnia</i> species (Crustacea, Cladocera) and the genetic characteristics of their populations based on allozyme studies in Lake Balaton, Hungary.....	79
SZŰTS, T.: Illustrations and redescriptions of Simon's little known salticid taxa from West-Africa (Araneae: Salticidae)	85
VÁRI, Á. & SZÖVÉNYI, G.: The habitat requirements of <i>Poecilimon brunneri</i> Frivaldszky, 1867 (Orthoptera: Phaneropteridae) and its Hungarian occurrence	97
<i>Communicatio Brevis</i>	
CSUZDI, CS. & POP, V. V.: Remarks on the species <i>Dendrobaena jeanneli</i> Pop, 1948 and its proposed synonymy with <i>Octolasion cyaneum</i> (Savigny, 1826) (Oligochaeta: Lumbricidae)	103